Current Neuropharmacology, 2022, 20, 269-271

COMMENTARY



Gut Serotonin as a General Membrane Permeability Regulator



Noémi Császár^{1,2} and Istvan Bókkon^{2,3,*}

¹National University of Public Services, Budapest, Hungary; ²Psychosomatic Outpatient Clinics, Budapest, Hungary; ³Vision Research Institute, Neuroscience and Consciousness Research Department, Lowell, MA, USA

ARTICLE HISTORY

Received: August 02, 2021 Revised: August 15, 2021 Accepted: August 15, 2021

10.2174/1570159X19666210921100542



1. COMMUNICATION BETWEEN THE GASTROINTESTINAL SYSTEM AND THE CENTRAL NERVOUS SYSTEM

The enteric nervous system (ENS) evolved before the central nervous system (CNS) and has

been considered as the "second brain" that can operate independently of the brain and spinal cord [1]. ENS may perform implicit learning and memorization, working like a little brain in the gut [2]. Perturbed gut microbiota homeostasis (dysbiosis) impairs bidirectional communication between the gastrointestinal (GI) system and the CNS (termed as the gut-brain axis) that is associated with neurodegenerative and neurological diseases [3-5].

2. SEROTONIN PRODUCED BY ENTERO-CHROMAFFIN CELLS IN THE GUT

Serotonin (5-HT) is an evolutionarily ancient molecule that is present in a wide range of species, from nematodes to humans [6]. 5-HT has key roles in numerous normal physiological conditions and pathological processes [7-10]. 5-HT is well-known as a neurotransmitter that regulates the neural activity and a wide range of neuropsychological processes [10]. Only a small fraction of 5-HT is manufactured in the brain since about 95% of the body's 5-HT is produced by enterochromaffin cells (ECs) in the gut that acts as a hormone with autocrine, paracrine, and endocrine functions [11]. 5-HT released from ECs can mediate diverse gastrointestinal functions like peristalsis, secretion, vasodilation, and perception of pain or nausea by means of activation of 5-HT receptors on intrinsic and extrinsic afferent nerve fibers that are found in the lamina propria [12]. 5-HT from the ECs is also picked up, to a large degree, by platelets and stored in their dense bodies and distributed throughout the body as a hormone and released upon their activation [13].

Dysbiosis has been linked to numerous chronic diseases like cardiovascular disease, obesity, diabetes, urinary stone disease, asthma, and inflammatory bowel disease, among others [14]. Dysbiosis that perturbs 5-HT levels of platelets and plasma [15, 16] is also associated with autism spectrum

disorder (APD), depression, anxiety, posttraumatic stress disorder (PTSD), pain, migraine, fibromyalgia, epilepsy, Parkinson's and Alzheimer's diseases, among others [17-22]. Psychological, environmental, and physical stressors also perturb the ENS and the gut microbial community that are also linked to multiple GI disorders and diverse diseases [23].

3. PLATELET AS CIRCULATING MIRRORS OF NEURONS

Almost one trillion platelets in the blood can work as immune cells and are essential mediators of hemostasis, thrombin generation, homeostasis, inflammation, and immune response [24-26]. In addition, platelets are increasingly considered a bridge between mental, immunological, and coagulation-related diseases [27]. Schizophrenia, depression, anxiety disorders, Parkinson's and Alzheimer's diseases, among others, are associated with platelet dysfunction [27]. Furthermore, platelets mimic the stable synaptic structure between neurons and mirror some features of neurons regarding protein expression [27-29]. It was proposed that platelets could be considered as circulating mirrors of neurons [28]. Both neurons and platelets have common proteins like reelin, amyloid peptides, Amyloid-beta precursor protein (APP), and Brain-derived neurotrophic factor (BDNF). Platelets contain cytokines (IL-1α, IL-1β, IL-4, TGFβ1), chemokines (CXCL4, CCL3), and neurotransmitters (serotonin, dopamine, epinephrine, histamine, gamma-aminobutyric acid (GABA)) [29, 30]. Activated platelets can also synthesize pro-inflammatory mediators such as Platelet-activating factor (PAF), Prostaglandins (PGs), and thromboxanes. Both platelets and neurons have similar secretory vesicles that contain 5-HT, dopamine, epinephrine, glutamate, GABA, calcium, adenosine 5'-diphosphate (ADP), and Adenosine 5'triphosphate (ATP), which are released from activated platelets or neurons following an action potential [27]. Serotonin released by activated platelets and Platelet-activating factor (PAF) have key functions in the regulation of sterile neuroinflammation, hemorrhage, and neuronal plasticity after traumatic brain injury [31].

4. INTESTINAL SEROTONIN AS A HORMONE-LIKE REGULATORY SIGNAL OF MEMBRANE PERMEA-BILITY

Recently, we proposed [32] that serotonin - produced by ECs - may work as a continuous hormone-like regulatory signal that influences membrane permeability in host organs

^{*}Address correspondence to this author at the Psychosomatic Outpatient Clinics, Budapest, Hungary; E-mail: bokkoni@yahoo.com

and tissues, including the brain. This signal by 5-HT is dependent on the intestine's actual health condition. Perturbed 5-HT biosynthesis in the gut may cause alterations in various platelet-dependent signal processes, including changes in vascular (membrane) permeability throughout the whole body as well as in the blood-brain barrier (BBB) [33, 34]. 5-HT induced changes in cellular permeability are able to affect many membrane-associated signaling processes. In addition, the gut microbiome influences glucose homeostasis of the whole body through gut-derived serotonin [35]. Furthermore, 5-HT has strong antioxidant and free radical scavenging ability, and its circulating levels are associated with a decrease in the plasma antioxidant capacity [36, 37]. Our concept may present an important role in which gut dysbiosis and 5-HT produced by ECs can considerably contribute to the development of a wide range of human diseases, including neurodevelopmental and neuropsychiatric disorders [32].

CONCLUSION

Cellular membrane signal processes have central roles in the regulation of cell functions and intra- and intercellular communication [38, 39]. Serotonin, produced by ECs and distributed by platelets, may work as a continuous hormone-like signal that can regulate membrane permeability in host organs and tissues, including the brain [32]. Understanding these processes could open up new opportunities in neuro(pharmacology) research. In addition, because serotonin transporters (SERTs) are present in the BBB, it suggests that 5-HT from the gut conveyed by platelets not only regulates the membrane permeability of the BBB, but 5-HT can also enter into the extracellular fluid of the CNS [40]. This 5-HT (from gut) may act by diffusion mechanism and may have a wide range of effects on the CNS via modulation of various signal and neurotransmission processes [41].

CONSENT FOR PUBLICATION

Not applicable.

FUNDING

None.

CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

ACKNOWLEDGEMENTS

The author, István Bókkon, gratefully acknowledge the assistance of SXM ExperiMental Ltd, Hungary, Budapest.

REFERENCES

- [1] Spencer, N.J.; Hibberd, T.J.; Travis, L.; Wiklendt, L.; Costa, M.; Hu, H.; Brookes, S.J.; Wattchow, D.A.; Dinning, P.G.; Keating, D.J.; Sorensen, J. Identification of a rhythmic firing pattern in the enteric nervous system that generates rhythmic electrical activity in smooth muscle. *J. Neurosci.*, 2018, 38(24), 5507-5522. http://dx.doi.org/10.1523/JNEUROSCI.3489-17.2018 PMID: 20807010
- [2] Schemann, M.; Frieling, T.; Enck, P. To learn, to remember, to forget-How smart is the gut? *Acta Physiol. (Oxf.)*, 2020, 228(1), e13296.

- http://dx.doi.org/10.1111/apha.13296 PMID: 31063665
- [3] Ma, Q.; Xing, C.; Long, W.; Wang, H.Y.; Liu, Q.; Wang, R.F. Impact of microbiota on central nervous system and neurological diseases: the gut-brain axis. *J. Neuroinflammation*, 2019, 16(1), 53. http://dx.doi.org/10.1186/s12974-019-1434-3 PMID: 30823925
- [4] Roy Sarkar, S.; Banerjee, S. Gut microbiota in neurodegenerative disorders. J. Neuroimmunol., 2019, 328, 98-104. http://dx.doi.org/10.1016/j.jneuroim.2019.01.004 PMID: 30658292
- [5] Singh, A.; Dawson, T.M.; Kulkarni, S. Neurodegenerative disorders and gut-brain interactions. *J. Clin. Invest.*, 2021, 131(13), e143775.
 http://dx.doi.org/10.1172/JCI143775 PMID: 34196307
- [6] Fidalgo, S.; Ivanov, D.K.; Wood, S.H. Serotonin: from top to bottom. *Biogerontology*, 2013, 14(1), 21-45. http://dx.doi.org/10.1007/s10522-012-9406-3 PMID: 23100172
- [7] Martin, A.M.; Young, R.L.; Leong, L.; Rogers, G.B.; Spencer, N.J.; Jessup, C.F.; Keating, D.J. The diverse metabolic roles of peripheral serotonin. *Endocrinology*, 2017, 158(5), 1049-1063. http://dx.doi.org/10.1210/en.2016-1839 PMID: 28323941
- [8] Kanova, M.; Kohout, P. Serotonin-Its synthesis and roles in the healthy and the critically Ill. Int. J. Mol. Sci., 2021, 22(9), 4837. http://dx.doi.org/10.3390/ijms22094837 PMID: 34063611
- [9] Filip, M.; Bader, M. Overview on 5-HT receptors and their role in physiology and pathology of the central nervous system. *Pharma*col. Rep., 2009, 61(5), 761-777. http://dx.doi.org/10.1016/S1734-1140(09)70132-X PMID: 19903999
- [10] Berger, M.; Gray, J.A.; Roth, B.L. The expanded biology of serotonin. *Annu. Rev. Med.*, **2009**, *60*, 355-366. http://dx.doi.org/10.1146/annurev.med.60.042307.110802 PMID: 19630576
- [11] Terry, N.; Margolis, K.G. Serotonergic mechanisms regulating the GI tract: Experimental evidence and therapeutic relevance. *Handb*. *Exp. Pharmacol.*, 2017, 239, 319-342. http://dx.doi.org/10.1007/164_2016_103 PMID: 28035530
- [12] Mawe, G.M.; Hoffman, J.M. Serotonin signalling in the gut-functions, dysfunctions and therapeutic targets. *Nat. Rev. Gastroenterol. Hepatol.*, 2013, 10(8), 473-486. http://dx.doi.org/10.1038/nrgastro.2013.105 PMID: 23797870
- [13] Lund, M.L.; Egerod, K.L.; Engelstoft, M.S.; Dmytriyeva, O.; The-odorsson, E.; Patel, B.A.; Schwartz, T.W. Enterochromaffin 5-HT cells A major target for GLP-1 and gut microbial metabolites. *Mol. Metab.*, 2018, 11, 70-83. http://dx.doi.org/10.1016/j.molmet.2018.03.004 PMID: 29576437
- [14] Wilkins, L.J.; Monga, M.; Miller, A.W. Defining dysbiosis for a cluster of chronic diseases. Sci. Rep., 2019, 9(1), 12918. http://dx.doi.org/10.1038/s41598-019-49452-y PMID: 31501492
- [15] Rogers, G.B.; Keating, D.J.; Young, R.L.; Wong, M.L.; Licinio, J.; Wesselingh, S. From gut dysbiosis to altered brain function and mental illness: mechanisms and pathways. *Mol. Psychiatry*, 2016, 21(6), 738-748. http://dx.doi.org/10.1038/mp.2016.50 PMID: 27090305
- [16] Yano, J.M.; Yu, K.; Donaldson, G.P.; Shastri, G.G.; Ann, P.; Ma, L.; Nagler, C.R.; Ismagilov, R.F.; Mazmanian, S.K.; Hsiao, E.Y. Indigenous bacteria from the gut microbiota regulate host serotonin biosynthesis. *Cell*, 2015, 161(2), 264-276. http://dx.doi.org/10.1016/j.cell.2015.02.047 PMID: 25860609
- [17] Padmakumar, M.; Van Raes, E.; Van Geet, C.; Freson, K. Blood platelet research in autism spectrum disorders: In search of biomarkers. *Res. Pract. Thromb. Haemost.*, 2019, 3(4), 566-577. http://dx.doi.org/10.1002/rth2.12239 PMID: 31624776
- [18] Murugesan, A.; Rani, M.R.S.; Hampson, J.; Zonjy, B.; Lacuey, N.; Faingold, C.L.; Friedman, D.; Devinsky, O.; Sainju, R.K.; Schuele, S.; Diehl, B.; Nei, M.; Harper, R.M.; Bateman, L.M.; Richerson, G.; Lhatoo, S.D. Serum serotonin levels in patients with epileptic seizures. *Epilepsia*, 2018, 59(6), e91-e97. http://dx.doi.org/10.1111/epi.14198 PMID: 29771456
- [19] Ernberg, M.; Voog, U.; Alstergren, P.; Lundeberg, T.; Kopp, S. Plasma and serum serotonin levels and their relationship to orofacial pain and anxiety in fibromyalgia. *J. Orofac. Pain*, 2000, 14(1), 37-46.
 PMID: 11203736
- [20] Tong, Q.; Zhang, L.; Yuan, Y.; Jiang, S.; Zhang, R.; Xu, Q.; Ding, J.; Li, D.; Zhou, X.; Zhang, K. Reduced plasma serotonin and 5-

- hydroxyindoleacetic acid levels in Parkinson's disease are associated with nonmotor symptoms. *Parkinsonism Relat. Disord.*, **2015**, *21*(8), 882-887.
- http://dx.doi.org/10.1016/j.parkreldis.2015.05.016 PMID: 26028271
- [21] Peitl, V.; Getaldić-Švarc, B.; Karlović, D. Platelet serotonin concentration is associated with illness duration in schizophrenia and chronological age in depression. *Psychiatry Investig.*, 2020, 17(6), 579-586. http://dx.doi.org/10.30773/pi.2020.0033 PMID: 32492767
- [22] Spivak, B.; Vered, Y.; Graff, E.; Blum, I.; Mester, R.; Weizman, A. Low platelet-poor plasma concentrations of serotonin in patients with combat-related posttraumatic stress disorder. *Biol. Psychiatry*, 1999, 45(7), 840-845. http://dx.doi.org/10.1016/S0006-3223(98)00231-5 PMID: 10202571
- [23] Karl, J.P.; Hatch, A.M.; Arcidiacono, S.M.; Pearce, S.C.; Pantoja-Feliciano, I.G.; Doherty, L.A.; Soares, J.W. Effects of psychological, environmental and physical stressors on the gut microbiota. Front. Microbiol., 2018, 9, 2013. http://dx.doi.org/10.3389/fmicb.2018.02013 PMID: 30258412
- [24] Chen, Y.; Zhong, H.; Zhao, Y.; Luo, X.; Gao, W. Role of platelet biomarkers in inflammatory response. *Biomark. Res.*, 2020, 8, 28. http://dx.doi.org/10.1186/s40364-020-00207-2 PMID: 32774856
- [25] Cognasse, F.; Laradi, S.; Berthelot, P.; Bourlet, T.; Marotte, H.; Mismetti, P.; Garraud, O.; Hamzeh-Cognasse, H. Platelet inflammatory response to stress. *Front. Immunol.*, 2019, 10, 1478. http://dx.doi.org/10.3389/fimmu.2019.01478 PMID: 31316518
- [26] Speth, C.; Löffler, J.; Krappmann, S.; Lass-Flörl, C.; Rambach, G. Platelets as immune cells in infectious diseases. *Future Microbiol.*, 2013, 8(11), 1431-1451. http://dx.doi.org/10.2217/fmb.13.104 PMID: 24199802
- [27] Izzi, B.; Tirozzi, A.; Cerletti, C.; Donati, M.B.; de Gaetano, G.; Hoylaerts, M.F.; Iacoviello, L.; Gialluisi, A. Beyond haemostasis and thrombosis: platelets in depression and its co-morbidities. *Int. J. Mol. Sci.*, 2020, 21(22), 8817. http://dx.doi.org/10.3390/ijms21228817 PMID: 33233416
- [28] Canobbio, I. Blood platelets: Circulating mirrors of neurons? Res. Pract. Thromb. Haemost., 2019, 3(4), 564-565. http://dx.doi.org/10.1002/rth2.12254 PMID: 31624775
- [29] Ponomarev, E.D. Fresh evidence for platelets as neuronal and innate immune cells: Their role in the activation, differentiation, and deactivation of Th1, Th17, and Tregs during tissue inflammation. Front. Immunol., 2018, 9, 406. http://dx.doi.org/10.3389/fimmu.2018.00406 PMID: 29599771
- [30] Pavlovic, V.; Ciric, M.; Jovanovic, V.; Trandafilovic, M.; Stojanovic, P. Platelet-rich fibrin: Basics of biological actions and protocol modifications. *Open Med. (Wars.)*, 2021, 16(1), 446-454. http://dx.doi.org/10.1515/med-2021-0259 PMID: 33778163
- [31] Dukhinova, M.; Kuznetsova, I.; Kopeikina, E.; Veniaminova, E.; Yung, A.W.Y.; Veremeyko, T.; Levchuk, K.; Barteneva, N.S.; Wing-Ho, K.K.; Yung, W.H.; Liu, J.Y.H.; Rudd, J.; Yau, S.S.Y.; Anthony, D.C.; Strekalova, T.; Ponomarev, E.D. Platelets mediate

- protective neuroinflammation and promote neuronal plasticity at the site of neuronal injury. *Brain Behav. Immun.*, **2018**, *74*, 7-27. http://dx.doi.org/10.1016/j.bbi.2018.09.009 PMID: 30217533
- [32] Szőke, H.; Kovács, Z.; Bókkon, I.; Vagedes, J.; Szabó, A.E.; Hegyi, G.; Sterner, M.G.; Kiss, Á.; Kapócs, G. Gut dysbiosis and serotonin: intestinal 5-HT as a ubiquitous membrane permeability regulator in host tissues, organs, and the brain. *Rev. Neurosci.*, 2020, 31(4), 415-425. http://dx.doi.org/10.1515/revneuro-2019-0095 PMID: 32007948
- [33] Sharma, H.S.; Olsson, Y.; Dey, P.K. Changes in blood-brain barrier and cerebral blood flow following elevation of circulating serotonin level in anesthetized rats. *Brain Res.*, **1990**, *517*(1-2), 215-223. http://dx.doi.org/10.1016/0006-8993(90)91029-G PMID: 2375992
- [34] Sharma, H.S.; Westman, J.; Navarro, J.C.; Dey, P.K.; Nyberg, F. Probable involvement of serotonin in the increased permeability of the blood-brain barrier by forced swimming. An experimental study using Evans blue and 131I-sodium tracers in the rat. *Behav. Brain Res.*, 1995, 72(1-2), 189-196. http://dx.doi.org/10.1016/0166-4328(96)00170-2 PMID: 8788871
- [35] Martin, A.M.; Yabut, J.M.; Choo, J.M.; Page, A.J.; Sun, E.W.; Jessup, C.F.; Wesselingh, S.L.; Khan, W.I.; Rogers, G.B.; Steinberg, G.R.; Keating, D.J. The gut microbiome regulates host glucose homeostasis via peripheral serotonin. Proc. Natl. Acad. Sci. USA, 2019, 116(40), 19802-19804. http://dx.doi.org/10.1073/pnas.1909311116 PMID: 31527237
- [36] Azouzi, S.; Santuz, H.; Morandat, S.; Pereira, C.; Côté, F.; Hermine, O.; El Kirat, K.; Colin, Y.; Le Van Kim, C.; Etchebest, C.; Amireault, P. Antioxidant and membrane binding properties of serotonin protect lipids from oxidation. *Biophys. J.*, 2017, 112(9), 1863-1873. http://dx.doi.org/10.1016/j.bpj.2017.03.037 PMID: 28494957
- [37] Amireault, P.; Bayard, E.; Launay, J.M.; Sibon, D.; Le Van Kim, C.; Colin, Y.; Dy, M.; Hermine, O.; Côté, F. Serotonin is a key factor for mouse red blood cell survival. *PLoS One*, 2013, 8(12), e83010. http://dx.doi.org/10.1371/journal.pone.0083010 PMID: 24358245
- [38] Yu, Y.; Li, M.; Yu, Y. Tracking Single Molecules in Biomembranes: Is Seeing Always Believing? ACS Nano, 2019, 13(10), 10860-10868. http://dx.doi.org/10.1021/acsnano.9b07445 PMID: 31589406
- [39] Grecco, H.E.; Schmick, M.; Bastiaens, P.I. Signaling from the living plasma membrane. Cell, 2011, 144(6), 897-909. http://dx.doi.org/10.1016/j.cell.2011.01.029 PMID: 21414482
- [40] Young, L.W.; Darios, E.S.; Watts, S.W. An immunohistochemical analysis of SERT in the blood-brain barrier of the male rat brain. *Histochem. Cell Biol.*, 2015, 144(4), 321-329. http://dx.doi.org/10.1007/s00418-015-1343-1 PMID: 26223876
- [41] Fuxe, K.; Borroto-Escuela, D.O. Volume transmission and receptor-receptor interactions in heteroreceptor complexes: understanding the role of new concepts for brain communication. *Neural Regen. Res.*, 2016, 11(8), 1220-1223. http://dx.doi.org/10.4103/1673-5374.189168 PMID: 27651759